



*This document is part of a collection of [Ecological Integrity Assessments](#) addressing 67 of Washington's 99 [Ecological Systems](#). These documents were prepared by the Washington Natural Heritage Program with funding provided by the Washington Department of Fish and Wildlife.*

---

## **Ecological Integrity Assessment: Rocky Mountain Subalpine-Montane Riparian Woodland**

### **Ecological Summary**

The Rocky Mountain Subalpine-Montane Riparian Woodland system is comprised of seasonally flooded forests and woodlands found at montane to subalpine elevations of the Rocky Mountain cordillera, from southern New Mexico north into Montana, and west into the Intermountain region and the Colorado Plateau. It occurs throughout the interior of British Columbia and the eastern slopes of the Cascade Mountains. In Washington, the system occurs at high elevations within dry and cold portions of the east Cascades, throughout the Okanogan Highlands, and in the Northern Rocky Mountains. Snowmelt moisture may create shallow water tables or seeps for a portion of the growing season. In Washington, stands typically occur at elevations between 2,000 – 7,000 feet (Kovalchik and Clausnitzer 2004). This system most commonly occurs in V-shaped, narrow valleys and canyons (where there is cold-air drainage). Less frequently, occurrences are found in moderate-wide valley bottoms on large floodplains along broad, meandering rivers, and on pond or lake margins.

Riparian woodland development is driven mostly by the magnitude and frequency of flooding, valley type, and stand replacing disturbances such as crown-fire, disease, windthrow, or clearcutting by humans or beaver. Valley type may be the most important variable, as riparian woodlands are mostly found in V-shaped, steep valleys with many large boulders and coarse soils. The forest vegetation in these environments is often very similar to the adjacent uplands (Baker 1987, Kovalchik and Clausnitzer 2004). Disturbances may create gaps in the canopy and allows pioneer species, such as aspen, or shrubs to establish. Less steep and wider valleys can lead to shrubland or woodland development. Flooding inundates vegetation, can physically dislodge seedlings/saplings, and alter channel morphology through erosion and deposition of sediment. Infrequent, high-powered floods determine large geomorphic patterns that persist on the landscape for hundreds to thousands of years (Hubert 2004). Floods of intermediate frequency and power produce floodplain landforms which persist for tens to hundreds of years as well as reset succession to early seral vegetation types (LANDIRE 2005; Hubert 2004). High frequency low-powered floods which occur nearly annually determine short-term patterns such as seed germination and seedling survival (Hubert 2004).

Narrow and steep (i.e. confined) occurrences have minimal to no floodplain development whereas less steep and wider valley bottoms (i.e., unconfined) occurrences are often associated with substantial floodplain development (LANDFIRE 2005; Gregory et al. 1991). Floodplains associated with the latter are comprised of a complexity of geomorphic surfaces which support a diverse array of vegetation communities and are able to store and release water slowly

throughout the growing season (Hubert 2004). Confined streams typically have shallow soils with minimal alluvium and transport water downstream rapidly through step-pool channels armored by boulders, bedrock, and large woody debris (LANDFIRE 2005; Hubert 2004).

Beaver, are of minimal significance in confined riparian woodlands as the steep nature of the latter system and often lack of deciduous trees typically precludes beaver activity. However, beaver activity but can have an impact on hydrology and vegetation in unconfined occurrences and especially in those areas dominated by aspen (*Populus tremuloides*).

Conifer and aspen woodlands dominate the canopy of this system. In Washington, confined occurrences (mostly along Rosgen A and B channels) are dominated by *Abies lasiocarpa* and/or *Picea engelmannii* (Kovalchik and Clausnitzer 2004). In older stands, *Picea engelmannii* may dominate the canopy while *Abies lasiocarpa* forms multi-aged canopies in the understory (Kovalchik and Clausnitzer 2004). Both *Abies lasiocarpa* and/or *Picea engelmannii* may be reproducing in the understory. *Pinus contorta*, *Pseudotsuga menziesii*, and *Larix occidentalis* are common early seral species. Common understory shrubs in confined woodlands include *Alnus viridis* ssp. *sinuata*, *Lonicera involucrata*, *Oplopanax horridus*, *Rosa gymnocarpa*, *Rubus parviflorus*, *Cornus canadensis*, *Ledum glandulosum*, *Vaccinium scoparium*, and *V. cespitosum*. *Arnica latifolia*, *Clintonia uniflora*, *Galium trifidum*, *Polemonium pulcherrimum*, *Senecio triangularis*, *Maianthemum stellatum*, *Streptopus amplexifolius*, *Athyrium filix-femina*, and *Gymnocarpium dryopteris* are common herbaceous species (Kovalchik and Clausnitzer 2004). Unconfined occurrences (mostly Rosgen C and E channels) are most often dominated by a canopy of *Picea engelmannii* while *Populus tremuloides*, *Betula papyrifera*, and occasionally *Pinus contorta* occur as early seral species. Common shrubs include *Cornus sericea*, *Symphoricarpos albus*, *Cornus canadensis*, *Lonicera involucrata*, *Rubus parviflorus*, *Pachistima myrsinites*, *Salix* ssp. *Alnus incana*, *A. viridis* ssp. *sinuata*, and *Ribes lacustre*. Herbaceous species often found in unconfined occurrences include *Carex scopulorum* var. *prionophylla*, *C. disperma*, *Elymus glaucus*, *Aralia nudicaulis*, *Streptopus amplexifolius*, *Gymnocarpium dryopteris*, and *Equisetum* ssp. Riparian woodlands dominated by *Populus tremuloides* are less common than coniferous dominated sites, however they can be found along riparian zones along low to moderate gradient channels (mostly Rosgen C and B channels) and ephemeral draws or depressions (Kovalchik and Clausnitzer 2004). Moderately large *Populus tremuloides* individuals are found in mature stands. *Betula papyrifera* and *Pinus contorta* are occasionally found in these stands. Regenerating *Populus tremuloides* and occasionally *Betula papyrifera*, *Pseudotsuga menziesii*, or *Picea engelmannii* can be found in the understory. Shrub diversity can be high and include *Cornus sericea*, *Symphoricarpos albus*, *Alnus incana*, *Acer glabrum* var. *douglasii*, *Amelanchier alnifolia*, *Ribes lacustre*, *Rosa gymnocarpa*, *Rubus parviflorus*, and *Salix* ssp. Herbaceous species are sparse in stands with high shrub cover. However, species such as *Carex pellita*, *Calamagrostis canadensis*, *Deschampsia cespitosa*, *Angelica arguta*, *Fragaria virginiana* var. *platypetala*, *Petasites sagittatus*, *Maianthemum stellatum*, and *Equisetum arvense* are often found in these woodlands (Kovalchik and Clausnitzer 2004). Some stands in northeastern Washington are dominated by *Thuja plicata* and/or *Tsuga heterophylla* and represent an inland version of the North Pacific Lowland Riparian Forest and Shrubland system. Either the EIA associated with the latter or this system should be applicable to these riparian forests. *Thuja plicata* and *Tsuga heterophylla* also occur along with *Oplopanax horridus* and

*Lysichiton americanus* on saturated soils in depressions or seeps. Such sites would be classified as the Northern Rocky Mountain Conifer Swamp.

The moisture associated with riparian areas promotes lower fire frequency compared with adjacent uplands. Stand replacement fires are rare but may occur when replacement fires occur in adjacent uplands (Fire regime III; average fire frequency of 100 years; LANDFIRE 2005). More frequent surface fires (~ every 50 years) can affect shrub patches through a combination of replacement fire from uplands and occasional native burning (LANDFIRE 2005). Following stand replacement fires deciduous woody species (e.g., *Populus tremuloides*, *Salix* spp., etc.) can be top-killed but generally resprout within a short period. Post-fire establishment of conifers occurs from seed. Wet meadows seldom burn and when they do, they typically recover within a single growing season (LANDFIRE 2005).

### *Stressors*

The stressors described below are those primarily associated with the loss of extent and degradation of the ecological integrity of existing occurrences. The stressors are the cause of the system shifting away from its natural range of variability. In other words, type, intensity, and duration of these stressors is what moves a system's ecological integrity rank away from the expected, natural condition (e.g. A rank) toward degraded integrity ranks (i.e. B, C, or D).

Historic and contemporary land use practices have impacted hydrologic, geomorphic, and biotic structure and function of riparian areas in eastern Washington. Human land uses both within the riparian area as well as in adjacent and upland areas have fragmented many riparian reaches which has reduced connectivity between riparian patches and riparian and upland areas. This can adversely affect the movement of surface/groundwater, nutrients, and dispersal of plants and animals. Roads, bridges, and development can also fragment both riparian and upland areas. Intensive grazing and recreation can also create barriers to ecological processes.

Reservoirs, water diversions, ditches, roads, and human land uses in the contributing watershed can have a substantial impact on the hydrology as well as biotic integrity of riparian woodlands (Woods 2001; Kattelman and Embury 1996; Poff et al. 1997; Baker 1987). All these stressors can induce downstream erosion and channelization, reduce changes in channel morphology, reduce base and/or peak flows, lower water tables in floodplains, and reduce sediment deposition in the floodplain (Poff et al. 1997). Vegetation responds to these changes by shifting from wetland and riparian dependent species to more mesic and xeric species typical of adjacent uplands (typical of herbaceous species) and/or encroaching into the stream channel. Although already narrow, floodplain width and the abundance and spatial distribution of various patch types also typically decline.

Livestock grazing is not a significant threat in confined riparian woodlands. However, in unconfined reaches, excessive livestock or native ungulate use can impact riparian woodlands by altering nutrient concentrations and cycles, changing surface and subsurface water movement and infiltration, shifting species composition, and reducing regeneration of woody species (Kauffman and Krueger 1984; Elmore and Kauffman 1984; Weixelman et al. 1997; Flenniken et al. 2001; Kauffman et al. 2004).

Management effects on woody riparian vegetation can be obvious, e.g., removal of vegetation by dam construction, roads, logging, or they can be subtle, e.g., removing beavers from a watershed, removing large woody debris, or construction of a weir dam for fish habitat. Non-native plants or animals, which can have wide-ranging impacts, also tend to increase with these stressors.

All of these stressors have resulted in many riparian areas being incised, supporting altered riparian plant communities, as well as numerous non-native species. This system has also decreased in extent due to agricultural development, roads, dams and other flood-control activities.

*Conceptual Ecological Model*

The general relationships among the key ecological attributes associated with this system are presented in Figure 1.




Figure 1. Conceptual Ecological Model for Rocky Mountain Subalpine-Montane Riparian Woodland.

**Ecological Integrity Assessments**

The assessment of ecological integrity can be done at three levels of intensity depending on the purpose and design of the data collection effort. The three-level approach is intended to provide increasing accuracy of ecological integrity assessment, recognizing that not all conservation and management decisions need equal levels of accuracy. The three-level approach also allows users to choose their assessment based in part on the level of classification that is available or targeted. If classification is limited to the level of forests vs. wetlands vs. grasslands, the use of remote sensing metrics may be sufficient. If very specific, fine-scale forest, wetland, and grassland types are the classification target then one has the flexibility to decide to use any of the three levels, depending on the need of the assessment. In other words, there is no presumption that a fine-level of classification requires a fine-level of ecological integrity assessment.

Because the purpose is the same for all three levels of assessment (to measure the status of ecological integrity of a site) it is important that the Level 1 assessment use the same kinds of metrics and major attributes as used at Levels 2 and 3. Level 1 assessments rely almost entirely on Geographic Information Systems (GIS) and remote sensing data to obtain information about landscape integrity and the distribution and abundance of ecological types in the landscape or watershed. Level 2 assessments use relatively rapid field-based metrics that are a combination of qualitative and narrative-based rating with quantitative or semi-quantitative ratings. Field observations are required for many metrics, and observations will typically require professional expertise and judgment. Level 3 assessments require more rigorous, intensive field-based methods and metrics that provide higher-resolution information on the integrity of occurrences. They often use quantitative, plot-based protocols coupled with a sampling design to provide data for detailed metrics.

Although the three levels can be integrated into a monitoring framework, each level is developed as a stand-alone method for assessing ecological integrity. **When conducting an ecological integrity assessment, one need only complete a single level that is appropriate to the study at hand.** Typically only one level may be needed, desirable, or cost effective. But for this reason it is very important that each level provide a comparable approach to assessing integrity, else the ratings and ranks will not achieve comparable information if multiple levels are used.

### **Level 1 EIA**

A generalized Level 1 EIA is provided in Rocchio and Crawford (2009). Please refer to that document for the list of metrics applicable to this ecological system.

## Level 2 EIA

The following table displays the metrics chosen to measure most of the key ecological attributes in the conceptual ecological model above. The EIA is used to assess the ecological condition of an assessment area, which may be the same as the element occurrence or a subset of that occurrence based on abrupt changes in condition or on artificial boundaries such as management areas. **Unless otherwise noted, metric ratings apply to both Level 2 and Level 3 EIAs. The difference between the two is that a Level 3 EIA will use more intensive and precise methods to determine metric ratings.** To calculate ranks, each metric is ranked in the field according to the ranking categories listed below. Then, the rank and point total for each metric is entered into the EIA Scorecard and multiplied by the weight factor associated with each metric resulting in a metric 'score'. Metric scores within a key ecological attribute are then summed to arrive at a score (or rank). These are then tallied in the same way to arrive at an overall ecological integrity score.

Table 1. Rocky Mountain Subalpine-Montane Riparian Woodland Level 2 EIA.

Metric	Justification	Rank			
		A (5 pts.)	B (4 pts.)	C (3 pts.)	D (1 pts.)
<b>Rank Factor: LANDSCAPE CONTEXT</b>					
<b>Key Ecological Attribute: <i>Buffer</i></b>					
<b>Buffer Length</b>	The buffer can be important to biotic and abiotic aspects of the wetland. Buffer Width Slope Multiplier 5-14% -->1.3; 15-40%-->1.4; >40%-->1.5	Buffer is > 75 – 100% of occurrence perimeter.	Buffer is > 50 – 74% of occurrence perimeter.	Buffer is 25 – 49% of occurrence perimeter	Buffer is < 25% of occurrence perimeter.
<b>Buffer Width</b>		Average buffer width of occurrence is > 200 m, adjusted for slope.	Average buffer width is 100 – 199 m, after adjusting for slope.	Average buffer width is 50 – 99 m, after adjusting for slope.	Average buffer width is < 49 m, after adjusting for slope.
<b>Buffer Condition</b>		Abundant (>95%) cover native vegetation, little or no (<5%) cover of non-native plants, intact soils, AND little or no trash or refuse.	Substantial (75–95%) cover of native vegetation, low (5–25%) cover of non-native plants, intact or moderately disrupted soils; minor intensity of human visitation or recreation.	Moderate (25–50%) cover of non-native plants, moderate or extensive soil disruption; moderate intensity of human visitation or recreation.	Dominant (>50%) cover of non-native plants, barren ground, highly compacted or otherwise disrupted soils, moderate or greater intensity of human visitation or recreation, no buffer at all.
<b>Key Ecological Attribute: <i>Landscape Structure</i></b>					

<b>Watershed Connectivity</b>	The types of land cover/uses in the contributing watershed has a significant effect on ecological processes(May 2002)	Landscape of contributing watershed primarily natural land cover; no connectivity barriers and no regional flood control dams upstream; <5% of contributing watershed urban or agricultural land cover types; few to no recent (<20 years) clearcut (<10% of the landscape)	5-20% of contributing watershed urban or agricultural land cover types; connectivity mostly retained; heavily managed forest landscape with many tree plantations (<50% of watershed in recent clearcuts)	20-50% of contributing watershed urban or agricultural land cover types; limited connectivity; heavily managed forest landscape with many tree plantations (<50% of watershed in recent clearcuts)	>50% of contributing watershed urban or agricultural land cover types; limited connectivity largely disrupted; one or more regional flood control dams upstream.
<b>Landscape Fire Regime Condition</b>	In mixed severity fire landscapes, fire effects can be out of Natural Range of Variability (Dwire and Kauffman 2003).	FRCC 1 No departure from historic fire regime.	FRCC 2 Slight-moderate departure from historic fire regime.		FRCC 3 Severe departure from historic fire regime. Fire suppression is evident; Fuel laddering is severe and throughout much of stand.
<b>Landscape Condition Model Index</b>	The intensity and types of land uses in the surrounding landscape can affect ecological integrity.	Landscape Condition Model Index > 0.8		Landscape Condition Model Index 0.65 – 0.79	Landscape Condition Model Index < 0.65
<b>Rank Factor: CONDITION</b>					
<b>Key Ecological Attribute: <i>Vegetation Composition</i></b>					
<b>Relative Cover Native Plant Species</b>	Native species dominate this system; non-natives increase with human impacts.	Cover of native plants 95-100%.	Cover of native plants 80-95%.	Cover of native plants 50 to 79%.	Cover of native plants <50%.
<b>Absolute Cover of Invasive Species</b>	Invasive species can inflict a wide range of ecological impacts. Early detection is critical. <i>Phalaris arundinacea</i> , are examples.	None present.	Invasive species present, but sporadic (<3% cover).	Invasive species prevalent (3–10% absolute cover).	Invasive species abundant (>10% absolute cover).
<b>Relative Cover of Native Increasers</b>	Some stressors such as grazing can shift or homogenize native composition toward species tolerant of stressors.	Absent or incidental	<10% cover	10-20% cover	>20% cover

<p><b>Species Composition</b> Note: Once developed, the Floristic Quality Assessment index could be used here instead.</p>	<p>The overall composition of native species can shift when exposed to stressors.</p>		<p>Species diversity/abundance at or near reference standard conditions. Native species sensitive to anthropogenic degradation are present, functional groups indicative of anthropogenic disturbance (ruderal or “weedy” species) are absent to minor, and full range of diagnostic / indicator species are present.</p>	<p>Species diversity/abundance close to reference standard condition. Some native species reflective of past anthropogenic degradation present. Some indicator/ diagnostic species may be absent.</p>	<p>Species diversity/abundance is different from reference standard condition in, but still largely composed of native species characteristic of the type. This may include ruderal (“weedy”) species. Many indicator/diagnostic species may be absent.</p>	<p>Vegetation severely altered from reference standard. Expected strata are absent or dominated by ruderal (“weedy”) species, or comprised of planted stands of non-characteristic species, or unnaturally dominated by a single species. Most or all indicator/diagnostic species are absent.</p>	
<p><b>Key Ecological Attribute: <i>Vegetation Structure</i></b></p>							
<p><b>Canopy structure</b></p>	<p>Intact riparian areas should have a diversity of tree age classes.</p>		<p>Average tree cover generally &gt; 50%; mixed age. Mature conifers and/or aspen and birch present; Trees are of sufficient size to provide future LWD to stream or floodplain.</p>		<p>Somewhat homogeneous in density and age OR &lt;50% canopy cover</p>	<p>Canopy extremely homogeneous, sparse, or absent (&lt;10% cover).</p>	
<p><b>Regeneration of Woody Species</b></p>	<p>Regeneration of woody species is expected in riparian areas with intact hydrology</p>		<p>Saplings/seedlings of native woody species (cottonwood/willow) present in expected amount; Obvious regeneration.</p>	<p>Saplings/seedlings of native woody species (cottonwood/willow) present but less than expected; Some seedling/saplings present.</p>	<p>Saplings/seedlings of native woody species (cottonwood/willow) present but in low abundance; Little regeneration by native species.</p>	<p>No reproduction of native woody species</p>	
<p><b>Large Woody Debris</b> (NMFS 1996, Fox 2001)</p>		<p>Large woody debris (LWD) in the stream channel is very important for channel formation, fish habitat, habitat heterogeneity, and sediment/hydrological processes</p>	<p>Bank-full Width 0-3 m</p>	<p>(piece = &gt;10cm diameter and ? 2 m in length; key piece &gt;28 pieces/100m of channel length</p>		<p>15-28</p>	<p>&lt;15</p>
		<p>&gt;3-30 m</p>	<p>&gt;3-30 m</p>	<p>&gt;56 pieces/100m of channel length</p>		<p>25-56</p>	<p>&lt;25</p>
		<p>&gt;30-50 m</p>	<p>&gt;30-50 m</p>	<p>&gt;63 pieces/100m of channel length</p>		<p>22-63</p>	<p>&lt;22</p>
<p><b>Patch Diversity and Connectivity</b></p>		<p>When hydrological processes are intact, a diversity of seral patch and habitat types are present within this system. The patches are well connected without interruption from anthropogenic land cover/use.</p>		<p>Confined reaches may have low patch diversity but connectivity within the riparian reach is unfragmented;  Unconfined reaches have heterogeneous mix of well connected patch types. Mature conifer, mature deciduous (aspen or birch)/conifer mixed, or mature deciduous patches present along with early seral stands of trees, wetland shrub and emergent vegetation patches.</p>	<p>Connectivity of confined reaches is becoming fragmented;  Unconfined reaches have expected patch diversity present but connectivity between patches is becoming fragmented. OR less diversity than expected, especially of mature stands of trees in unconfined reaches.</p>	<p>Confined reaches are moderately fragmented;  Patch diversity in unconfined reaches is low and becoming homogeneous in unconfined reaches; few if any mature stands of trees present. Many patches isolated due to fragmentation within the riparian system.</p>	<p>Confined reaches are severely fragmented;  Mostly dominated by one patch type (unconfined reaches). No mature conifer or deciduous tree patches present. Patch is isolated due to fragmentation within the riparian system.</p>

<b>Key Ecological Attribute: <i>Hydrology</i></b>					
<b>Water Source</b>	Anthropogenic sources of water can have detrimental effects on the hydrological regime	Source is natural or naturally lacks water in the growing season. No indication of direct artificial water sources	Source is mostly natural, but site directly receives occasional or small amounts of inflow from anthropogenic sources	Source is primarily urban runoff, direct irrigation, pumped water, artificially impounded water, or other artificial hydrology	Water flow has been substantially diminished by human activity
<b>Channel Stability</b>	Alteration in hydrology or sediment loads or some onsite stressors can degrade channel stability	Natural channel; no evidence of severe aggradation or degradation;	Most of the channel has some aggradation or degradation, none of which is severe	Evidence of severe aggradation or degradation of most of the channel	Concrete, or artificially hardened, channels through most of the site
<b>Streambank Stability</b>	Stable streambanks are indicative of intact hydrological and sediment regimes. (Henshaw and Booth 2000)	Stable Perennial vegetation to waterline; no raw or undercut banks (some erosion on outside of banks normal); no recently exposed roots; no recent tree falls	Slightly Stable Perennial vegetation to waterline in most places; minor erosion and/or bank undercutting; recently exposed tree roots rare but present	Moderately Unstable Perennial vegetation to waterline sparse (mainly scoured or removed by lateral erosion); bank held in place by hard points (trees, boulders) and eroded back elsewhere; extensive erosion and bank undercutting; recently exposed tree roots and fine root hairs common	Completely Unstable No perennial vegetation to waterline; banks only held by hard points; severe erosion of both banks; recently exposed tree roots common; tree falls and/or severely undercut trees common
<b>Hydrological Connectivity (Riverine)</b>	Floodwater should have access to the floodplain. Stressors resulting in entrenchment affect hydrological connectivity	<b>LEVEL 2:</b> Completely connected to floodplain (backwater sloughs and channels)	Minimally disconnected from floodplain by dikes, elevated culverts, etc	Moderately disconnected from floodplain by dikes, elevated culverts, etc.	Extensively disconnected from floodplain by dikes, elevated culverts, etc.
		<b>LEVEL 3:</b> Unconfined: Entrenchment ratio > 4.0; Confined: Entrenchment ratio > 1.4	Unconfined: Entrenchment ratio 1.4 – 2.2; Confined: Entrenchment ratio 1.0 – 1.4	Unconfined: Entrenchment ratio < 1.4; Confined: Entrenchment ratio < 1.0	
<b>Key Ecological Attribute: <i>Physicochemical</i></b>					
<b>Soil Surface Condition</b>	Soil disturbance can result in erosion thereby negatively affecting many ecological processes	Bare soil areas are limited to naturally caused disturbances such as flood deposition or game trails	Some bare soil due to human causes but the extent and impact is minimal. The depth of disturbance is limited to only a few inches and does not show evidence of ponding or channeling water.	Bare soil areas due to human causes are common. There may be pugging due to livestock resulting in several inches of soil disturbance. ORVs or other machinery may have left some shallow ruts.	Bare soil areas substantially & contribute to altered hydrology or other long-lasting impacts. Deep ruts from ORVs or machinery may be present, or livestock pugging and/or trails are widespread. Water will be channeled or ponded.

<b>Water Quality</b>	Excess nutrients, sediments, or other pollutant have an adverse affect on natural water quality	No evidence of degraded water quality. Water is clear; no strong green tint or sheen.	Some negative water quality indicators are present, but limited to small and localized areas. Water may have a minimal greenish tint or cloudiness, or sheen.	Negative indicators or wetland species that respond to high nutrient levels are common. Water may have a moderate greenish tint, sheen or other turbidity with common algae.	Widespread evidence of negative indicators. Algae mats may be extensive. Water may have a strong greenish tint, sheen or turbidity. Bottom difficult to see during due to surface algal mats and other vegetation blocking light to the bottom.
<b>Rank Factor: SIZE</b>					
<b>Key Ecological Attribute: <i>Size</i></b>					
<b>Relative Size</b>	Indicates the proportion lost due to stressors.	Site is at or minimally reduced from natural extent (>95% remains)	Occurrence is only modestly reduced from its original natural extent (80-95% remains)	Occurrence is substantially reduced from its original natural extent (50-80% remains)	Occurrence is severely reduced from its original natural extent (<50% remains)
<b>Absolute Size</b>	Absolute size may be important for buffering impacts originating in the surrounding landscape	Streams with limited floodplain development, primarily braided channels, or extremely sinuous stable channels (Rosgen A, B, D, E, or F types)  > 5.0 linear miles (8 km)	3.0 to 5.0 linear miles (5-8 km)	1.0 to 3.0 miles linear miles (1.5 to 5 km)	< 1.0 linear miles (1.5 km)
		Meandering streams with well-developed floodplains and wide channels (mostly Rosgen C type)  >25 meander wavelengths or 50 point bars	10-25 meander wavelengths or 20-50 point bars	4-10 meander wavelengths or 8-20 point bars	<4 meander wavelengths or <8 point bars

### Level 3 EIA

Level 3 metrics would include more quantitative measures of the metrics listed above. In addition, the following metrics should be considered in a Level 3 EIA:

- Benthic invertebrate Index of Biotic Integrity (BIBI; WADOE 2003); Statewide data are maintained by WADOE:  
<http://www.ecy.wa.gov/apps/watersheds/streambio/regions/state.asp?symtype=1>
- Index of Hydrological Alteration (Richter et al. 1997)
- Specific water quality measures (e.g., the temperature, dissolved oxygen, pH, conductivity, turbidity of stream water)
- Pool Quality Index (May (2002); may need modification for Eastside riparian systems)
- Riffle Quality Index (May (2002); may need modification for Eastside riparian systems)

### Triggers or Management Assessment Points

Ecological triggers or conditions under which management activities need to be reassessed are shown in the table below. Since the Ecological Integrity rankings are based on hypothesized thresholds, they are used to indicate where triggers might occur. Specific details about how these triggers translate for each metric can be found by referencing the values or descriptions for the appropriate rank provided in the Table above.

Table 2. Triggers for Level 2 & 3 EIA

Key Ecological Attribute or Metric	Trigger	Action
Any metric (except Connectivity)	<ul style="list-style-type: none"> <li>▪ C rank</li> <li>▪ Shift from A to B rank</li> <li>▪ negative trend within the B rating (Level 3)</li> </ul>	<p>Level 2 triggers: conduct Level 3 assessment; make appropriate short-term management changes to ensure no further degradation</p> <p>Level 3 triggers: make appropriate management adjustments to ensure no additional degradation occurs. Continue monitoring using Level 3.</p>
Any Key Ecological Attribute	<ul style="list-style-type: none"> <li>▪ any metric has a C rank</li> <li>▪ &gt; ½ of all metrics are ranked B</li> <li>▪ negative trend within the B rating (Level 3)</li> </ul>	<p>Level 2 triggers: conduct Level 3 assessment; make appropriate short-term management changes to ensure no further degradation</p> <p>Level 3 triggers: make appropriate management adjustments to ensure no additional degradation occurs. Continue monitoring using Level 3.</p>

### Protocol for Integrating Metric Ranks

If desired, the user may wish to integrate the ratings of the individual metrics and produce an overall score for the three rank factor categories: (1) Landscape Context; (2) Condition; and (3) Size. These rank factor rankings can then be combined into an Overall Ecological Integrity Rank. This enables one to report scores or ranks from the various hierarchical scales of the assessment depending on which best meets the user’s objectives. Please see Table 5 in Rocchio and Crawford (2009) for specifics about the protocol for integrating or ‘rolling-up’ metric ratings.

Supporting documents for the EIAs can be found at:  
<http://www1.dnr.wa.gov/nhp/refdesk/communities/eia.html>

Documentation about Ecological Systems can be found at:  
[http://www1.dnr.wa.gov/nhp/refdesk/communities/ecol\\_systems.html](http://www1.dnr.wa.gov/nhp/refdesk/communities/ecol_systems.html)

## References

Baker, W.L. 1987. Recent Changes in the Riparian Vegetation of the Montane and Subalpine Zones of Western Colorado, U.S.A. PhD Dissertation. University of Wisconsin. Madison, WI.

Collins, J.N., E. Stein, and M. Sutula. 2008. California rapid assessment method (CRAM) for wetlands. Version 5.0.2. San Francisco Estuary Institute. San Francisco, California. Available online at: <http://www.cramwetlands.org/>

Crawford, R.C. and J. Kagan. 2001. Eastside Riparian-Wetlands. *In* Johnson, D.H. and T.A. O'Neil. 2001. Wildlife-Habitat Relationships in Oregon and Washington. Oregon State University Press, Corvallis, OR.

Dwire, K.A. and J.B. Kauffman. 2003. Fire and riparian ecosystems in landscapes of the western USA. *Forest Ecology and Management* 178 : 61–74

Elmore, W. and B. Kauffman. 1994. Riparian and Watershed Systems: Degradation and Restoration. *In*: Ecological implications of livestock herbivory in the west. Society of Range Mgmt. Denver, CO.

Flenniken, M., R.R. McEldowney, W.C. Leininger, G.W. Frasier, and M.J. Trlica. Hydrologic responses of a montane riparian ecosystem following cattle use. *Journal of Range Management* 54: 567-574.

Fox, M.J. 2001. A New Look at the Quantities and Volumes of Instream Wood in Forested Basins within Washington State. M.S. Thesis. College of Forest Resources, University of Washington. Seattle, WA. Online: <http://water.washington.edu/Theses/fox.html>

Gregory, S.V., F.J. Swanson, W.A. McKee, and K.W. Cummins. 1991. An Ecosystem Perspective of Riparian Zones. *BioScience* 41(8): 540-551.

Henshaw, P.C. and Booth, D.B., 2000. Natural Restabilization of Stream Channels in Urban Watersheds. *Journal of the American Water Resources Association*, 36(6): 1219-1236.

Hubert, W.A. 2004. Ecological Processes in Riverine Wetland Habitats. Pages 52-73 *in* M. C. McKinstry, W.A. Hubert, and S.H. Anderson, editors. *Wetland and Riparian Areas of the Intermountain West: Ecology and Management*. University of Texas Press, Austin, TX.

Kattelman, R. and M. Embury. 1996. Riparian Areas and Wetlands. *In* Sierra Nevada Ecosystem Project: Final Report to Congress, Vol. III, Assessment and Scientific Basis for

Management Options. Center for Water and Wildland Resources, University of California, Davis, CA.

Kauffman, J.B. and W.C. Krueger. 1984. Livestock Impacts on Riparian Ecosystems and Streamside Management Implications: A Review. *J. Range Manage.* 37(5):430-438.

Kauffman, J.B., A.S. Thorpe, and E.N.J. Brookshire. 2004. Livestock exclusion and belowground ecosystem responses in riparian meadows of eastern Oregon. *Ecological Applications* 14(6): 1671-1679.

Kovalchik, B.L. and R.R. Clausnitzer. 2004. Classification and Management of Aquatic, Riparian, and Wetland Sites on the National Forests of Eastern Washington. Series Description. Pacific Northwest Research Station, Forest Service, U.S. Department of Agriculture. In cooperation with: Pacific Northwest Region, Colville, Okanogan, and Wenatchee National Forests. General Technical Report PNW-GTR-593. Portland, OR.

LANDFIRE. 2005. Landfire Biophysical Setting Model for Rocky Mountain Subalpine/Upper Montane Riparian Systems. BpS1011600. Online: <http://www.landfire.gov/NationalProductDescriptions24.php>

Laubhan, M.K. 2004. Variation in Hydrology, Soils, and Vegetation of Natural Palustrine Wetlands Among Geologic Provinces. Pages 23-51 *in* M. C. McKinstry, W.A. Hubert, and S.H. Anderson, editors. *Wetland and Riparian Areas of the Intermountain West: Ecology and Management*. University of Texas Press, Austin, TX.

May, C.W. 2002. Measures of Ecological Integrity for Salmonid Streams on Department of Defense Facilities in the Pacific Northwest: Current Watershed Conditions and Management Recommendations. Technical Report APL-UWTR0104. Applied Physics Laboratory, University of Washington, Seattle, WA. Online: <http://www.serdp.org/Research/upload/CS-1154-FR-01.pdf>

NatureServe Explorer. 2007. Descriptions of Ecological Systems for the State of Washington. Data current as of October 06, 2007. NatureServe, Arlington, VA. [\[http://www.natureserve.org/explorer/index.htm\]](http://www.natureserve.org/explorer/index.htm)

Poff, N.L., J.D. Allan, M.B. Bain, J.R. Karr, K.L. Presegaard, B.D. Richter, R.E. Sparks, and J.C. Stromburg. 1997. The Natural Flow Regime: A Paradigm for River Conservation and Restoration. *BioScience* 47: 769-784.

Richter, B.D., J.V. Baumgartner, J. Powell, and D.P. Braun. 1997. A Method for Assessing Hydrologic Alteration within Ecosystems. *Conservation Biology* 10: 1163-1174.

Rink, L.P. and G.N. Kiladis. 1986. Geology, hydrology, climate, and soils of the Rocky Mountains. Pages 42-65 *in* J.T. Windell, B.E. Willard, D.J. Cooper, S.Q. Foster, C.F. Knud-Hansen, L.P. Rink, and G.N. Kiladis, editors. *An ecological characterization of Rocky Mountain montane and subalpine wetlands*. U.S. Fish and Wildlife Service, Biological Report 86.

Rocchio, F.J. and R.C. Crawford. 2009. Monitoring Desired Ecological Conditions on Washington State Wildlife Areas Using an Ecological Integrity Assessment Framework. Washington Natural Heritage Program, Washington Department of Natural Resources, Olympia, WA.

Rocchio, J. 2006. Rocky Mountain Subalpine-Montane Riparian Woodland Ecological Integrity Assessment. Report Prepared for NatureServe, Arlington, VA. Colorado Natural Heritage Program, Colorado State University. Fort Collins, CO. Online: [http://www.cnhp.colostate.edu/download/documents/2005/ecological\\_integrity/Rocky%20Mountain%20Subalpine-Montane%20Riparian%20Woodlands\\_EIA\\_Dec09\\_05.pdf](http://www.cnhp.colostate.edu/download/documents/2005/ecological_integrity/Rocky%20Mountain%20Subalpine-Montane%20Riparian%20Woodlands_EIA_Dec09_05.pdf)

Washington Department of Ecology (WADOE). 2003. Multi-metric Index Development for Biological Monitoring in Washington State Streams. Publication 03-03-035. Washington Department of Ecology, Olympia, WA. Online: <http://www.ecy.wa.gov/pubs/0303035.pdf>

Weixelman D.A., D.C. Zamudio, K.A. Zumudio, and R.T. Tausch. 1997. Classifying ecological types and evaluating site degradation. *Journal of Range Management* 50: 315-321.

Woods, S.W. 2001. Ecohydrology of subalpine wetlands in the Kawuneeche Valley, Rocky Mountain National Park, Colorado. PhD Dissertation. Department of Earth Sciences, Colorado State University, Fort Collins, CO.

Authorship: Joe Rocchio, Washington Natural Heritage Program  
February 22, 2011