

***Campanula lasiocarpa* Cham.**  
Alaska harebell  
Campanulaceae (Harebell Family)

**Status:** State Sensitive  
**Rank:** G5S2

**General Description:** Adapted from Hitchcock et al. (1959): This rhizomatous perennial plant has 1 to several lax stems up to 6 in. (15 cm) tall. The plant lacks hair except for the hypanthium, which is moderately to densely covered with long matted hairs and/or long, soft, fine trichomes and the petioles or proximal margins of the leaves that are fringed with conspicuous hairs. The leaves are evidently to sometimes obscurely serrate. The basal leaves are petiolate, with oblanceolate to elliptic or subrhombic blades that are  $\frac{1}{4}$  to 1 in. (6 to 25 mm) long. The few cauline leaves are more or less reduced. The flowers are solitary or rarely two. The calyx lobes are leaflike but narrow,  $\frac{1}{4}$  to  $\frac{3}{4}$  in. (5 to 18 mm) long, and commonly some or all of them have a few slender teeth. The blue corolla is  $\frac{3}{4}$  to 1  $\frac{1}{8}$  in. (18 to 30 mm) long, surpassing the style, and the broad lobes are much shorter than the tube. The anthers are  $\frac{1}{8}$  to  $\frac{1}{4}$  (3 to 5 mm) long. The capsule is subcylindric, not much enlarged upward, about  $\frac{1}{4}$  in. (1 cm) long, and opening near the summit.

**Identification Tips:** The four closely related *species of Campanula* (*C. lasiocarpa*, *C. piperi*, *C. scabrella*, and *C. parryi*), only *C. lasiocarpa* and *C. scabrella* overlap in geographic location. These two species can be easily distinguished by the nature of their hypanthium, corolla, and leaves. The hypanthium of *C. lasiocarpa* is moderately to densely covered with long, loosely spreading hairs and/or long, soft, fine trichomes. Its corolla is mostly  $\frac{3}{4}$  to 1  $\frac{1}{8}$  in. (18 to 30 mm) long with lobes that are much shorter than the tube, and leaves that are slightly to strongly toothed. The hypanthium of *C. scabrella* is hairless to slightly rough because of the presence of short, stiff hairs and is sometimes also covered with minute rough, stiff trichomes. Its corolla is mostly  $\frac{1}{4}$  to  $\frac{2}{3}$  in. (6 to 16 mm) long with lobes that are much shorter than the tube and its leaves are entire or nearly so.

**Phenology:** This species flowers from July to August.

**Range:** This species is found in the Cascade Range of Snohomish and King counties, Washington, the Selkirk Range of British Columbia, north to Alaska, throughout the Aleutian Islands to Kamchatka and south to Hokkaido, Japan.


**Habitat:** *Campanula lasiocarpa* prefers rock crevices in alpine zones, usually in unglaciated areas. In the Mt. Baker-Snoqualmie National Forest it has been seen in dry, rocky microsites within cool, wet subalpine areas. Populations in Washington have been found at an elevation of 2000 to 6840 ft (610 to 2085

***Campanula lasiocarpa***  
Alaska harebell



©1959 Reprinted by permission of the University of Washington Press.  
Illustration by Jeanne R. Janish

Known distribution of  
*Campanula lasiocarpa*  
in Washington



## ***Campanula lasiocarpa***

Alaska harebell



## ***Campanula lasiocarpa***

Alaska harebell

m). Associated species include yellow mountain heather (*Phyllodoce glanduliflora*), three-forked artemisia (*Artemisia trifurcata*), and golden fleabane (*Erigeron aureus*).

**Ecology:** Globally this species grows in alpine heaths and sandy tundra.

**State Status Comments:** The species is peripheral in Washington. It is known from fewer than ten recent occurrences in King and Snohomish counties and one historical site. Extant populations are quite small.

**Inventory Needs:** Additional inventory is needed. Rocky areas within alpine habitats in King and Snohomish Counties should be systematically surveyed for additional populations. Known populations should be revisited and updated habitat and population information should be collected.

**Threats and Management Concerns:** Existing threats may include grazing and trampling from goats and trampling from climbers.

### **References:**

- Hitchcock, C.L., A. Cronquist, M. Ownbey, J.W. Thompson. 1959. *Vascular Plants of the Pacific Northwest Part 4: Ericaceae Through Campanulaceae*. University of Washington Press, Seattle, WA. 510 pp.
- NatureServe. 2003. NatureServe Explorer: An online encyclopedia of life [web application]. Version 1.8. NatureServe, Arlington, Virginia. Available <http://www.natureserve.org/explorer>. (Accessed: September 19, 2003).
- Potash, L.L. 1991. *Sensitive Plants and Noxious Weeds of the Mt. Baker-Snoqualmie National Forest*. United States Department of Agriculture: Forest Service, Washington. 116 pp.